

intravenous injection, regional administration via blood or lymph supply, or directly to an affected site. Also contemplated are transdermal routes, such as, via a patch.

Determining the effectiveness of a compound *in vivo* may involve a variety of different criteria. Also, measuring toxicity and dose response can be performed in animals in a more meaningful fashion than in *in vitro* or *in cyto* assays.

### III. Chromatographic Separation Procedures

Any of a wide variety of chromatographic procedures may be employed according to the present invention. For example, thin layer chromatography (TLC), gas chromatography, high performance liquid chromatography (HPLC), paper chromatography, affinity chromatography or supercritical flow chromatography may be used to effect separation of various chemical species.

Partition chromatography is based on the theory that if two phases are in contact with one another, and if one or both phases constitute a solute, the solute will distribute itself between the two phases. Usually, partition chromatography employs a column, which is filled with a sorbent and a solvent. The solution containing the solute is layered on top of the column. The solvent is then passed through the column, continuously, which permits movement of the solute through the column material. The solute can then be collected based on its movement rate. The two most common types of partition chromatograph are paper chromatography and thin-layer chromatography (TLC); together these are called adsorption chromatography. In both cases, the matrix contains a bound liquid. Other examples of partition chromatography are gas-liquid and gel chromatography.

Paper chromatography is a variant of partition chromatography that is performed on cellulose columns in the form of a paper sheet. Cellulose contains a large amount of bound water even when extensively dried. Partitioning occurs between the bound water and the developing solvent. Frequently, the solvent used is water. Usually, very small volumes of the solution mixture to be separated is placed at top of the paper and allowed

to dry. Capillar action draws the solvent through the paper, dissolves the sample, and moves the components in the direction of flow. Paper chromatograms may be developed for either ascending or descending solvent flow. Two dimensional separations are permitted by changing the axis of migration 90° after the first run.

5 Thin layer chromatography (TLC) is very commonly used to separate lipids and, therefore, is considered a preferred embodiment of the present invention. TLC has the advantages of paper chromatography, but allows the use of any substance that can be finely divided and formed into a uniform layer. In TLC, the stationary phase is a layer of sorbent spread uniformly over the surface of a glass or plastic plate. The plates are  
10 usually made by forming a slurry of sorbent that is poured onto the surface of the gel after creating a well by placing tape at a selected height along the perimeter of the plate. After the sorbent dries, the tape is removed and the plate is treated just as paper in paper chromatography. The sample is applied and the plate is contacted with a solvent. Once the solvent has almost reached the end of the plate, the plate is removed and dried. Spots  
15 can then be identified by fluorescence, immunologic identification, counting of radioactivity, or by spraying varying reagents onto the surface to produce a color change.

In gas-liquid chromatography (GLC), the mobile phase is a gas and the stationary phase is a liquid adsorbed either to the inner surface of a tube or column or to a solid support. The liquid usually is applied as a solid dissolved in a volatile solvent such as  
20 ether. The sample, which may be any sample that can be volatized, is introduced as a liquid with an inert gas, such as helium, argon or nitrogen, and then heated. This gaseous mixture passes through the tubing. The vaporized compounds continually redistribute themselves between the gaseous mobile phase and the liquid stationary phase, according to their partition coefficients.

25 The advantage of GLC is in the separation of small molecules. Sensitivity and speed are quite good, with speeds that approach 1000 times that of standard liquid chromatography. By using a non-destructive detector, GLC can be used preparatively to

purify grams quantities of material. The principal use of GLC has been in the separation of alcohols, esters, fatty acids and amines.

Gel chromatography, or molecular sieve chromatography, is a special type of partition chromatography that is based on molecular size. The theory behind gel chromatography is that the column, which is prepared with tiny particles of an inert substance that contain small pores, separates larger molecules from smaller molecules as they pass through or around the pores, depending on their size. As long as the material of which the particles are made does not adsorb the molecules, the sole factor determining rate of flow is the size. Hence, molecules are eluted from the column in decreasing size, so long as the shape is relatively constant. Gel chromatography is unsurpassed for separating molecules of different size because separation is independent of all other factors such as pH, ionic strength, temperature, etc. There also is virtually no adsorption, less zone spreading and the elution volume is related in a simple matter to molecular weight.

The gel material for gel chromatography is a three-dimensional network whose structure is usually random. The gels consist of cross-linked polymers that are generally inert, do not bind or react with the material being analyzed, and are uncharged. The space filled within the gel is filled with liquid and this liquid occupies most of the gel volume. Common gels are dextran, agarose and polyacrylamide; they are used for aqueous solution.

High performance liquid chromatography (HPLC) is characterized by a very rapid separation with extraordinary resolution of peaks. This is achieved by the use of very fine particles and high pressure to maintain an adequate flow rate. Separation can be accomplished in a matter of minutes, or at most an hour. Moreover, only a very small volume of the sample is needed because the particles are so small and close-packed that the void volume is a very small fraction of the bed volume. Also, the concentration of the sample need not be very great because the bands are so narrow that there is very little dilution of the sample.